

Manuscript version: Author's Accepted Manuscript

The version presented in WRAP is the author's accepted manuscript and may differ from the published version or Version of Record.

Persistent WRAP URL:

<http://wrap.warwick.ac.uk/114461/>

How to cite:

Please refer to published version for the most recent bibliographic citation information. If a published version is known of, the repository item page linked to above, will contain details on accessing it.

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

© 2016 Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International <http://creativecommons.org/licenses/by-nc-nd/4.0/>.



Publisher's statement:

Please refer to the repository item page, publisher's statement section, for further information.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk.

Accepted Manuscript

Title: Zinc Chloride Aqueous Solution as a Solvent for Starch

Author: Meiyang Lin Xiaoqin Shang Peng Liu Fengwei Xie
Xiaodong Chen Yongyi Sun Junyan Wan



PII: S0144-8617(15)00854-1
DOI: <http://dx.doi.org/doi:10.1016/j.carbpol.2015.09.007>
Reference: CARP 10309

To appear in:

Received date: 20-5-2015
Revised date: 30-8-2015
Accepted date: 2-9-2015

Please cite this article as: Lin, M., Shang, X., Liu, P., Xie, F., Chen, X., Sun, Y., and Wan, J., Zinc Chloride Aqueous Solution as a Solvent for Starch, *Carbohydrate Polymers* (2015), <http://dx.doi.org/10.1016/j.carbpol.2015.09.007>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Highlights:

- ✓ ZnCl_2 aqueous solution (>29.6 wt.%) is a non-derivatizing solvent for starch
- ✓ Starch can be dissolved completely and regenerated easily in ZnCl_2 solution
- ✓ The suggested dissolution mechanism was the formation of a “zinc-starch complex”, not degradation by H^+ .

Zinc Chloride Aqueous Solution as a Solvent for Starch

Meiying Lin^a, Xiaoqin Shang^{a*}, Peng Liu^a, Fengwei Xie^b, Xiaodong Chen^a, Yongyi Sun^a, Junyan Wan^a

^a School of Chemistry and Chemical Engineering, Guangzhou University, Guangzhou 510006, China

^b School of Chemical Engineering, The University of Queensland, Brisbane, Qld 4072, Australia

* Corresponding author. Tel.: +86 20 13342885662; fax: +86 20 39366902; Email address: *hushanren@163.com* (X.Q. Shang)

Abstract:

It is important to obtain starch-based homogeneous systems for starch modification. Regarding this, an important key point is to find cheap, low-cost and low-toxicity solvents to allow complete dissolution of starch and its easy regeneration. This study reveals that a ZnCl_2 aqueous solution is a good non-derivatizing solvent for starch at 50 °C, and can completely dissolve starch granules. The possible formation of a “zinc-starch complex” might account for the dissolution; and the degradation of starch, which was caused by the H^+ in ZnCl_2 aqueous solution, could not contribute to full dissolution. From polarized light microscopic observation combined with the solution turbidity results, it was found that the lowest ZnCl_2 concentration for full dissolution was 29.6 wt.% at 50 °C, with the dissolving time being 4 h. Using Fourier-transform infrared (FTIR), solid state ^{13}C nuclear magnetic resonance (NMR), and X-ray diffraction (XRD), it was revealed that ZnCl_2 solution had no chemical reaction with starch glucosides, but only weakened starch hydrogen bonding and converted the crystalline regions to amorphous regions. In addition, as shown by intrinsic viscosity and thermogravimetric analysis (TGA), ZnCl_2 solution caused degradation of starch macromolecules, which was more serious with a higher concentration of ZnCl_2 solution.

Keywords: starch; ZnCl_2 aqueous solution; non-derivatizing solvent; dissolution; zinc-starch complex

Chemical compounds studied in this article:

Starch (PubChem CID: 24836924); Water (PubChem CID: 962); Zinc dichloride (PubChem CID: 5727)

39

40 **1. Introduction**

41 Starch is one of the most abundant carbohydrates in nature, and has been widely
42 used as a raw material in industry. Furthermore, native starch can be modified into
43 various derivatives to widen its applications. Modified starches such as starch sodium
44 succinate and starch phosphate ester can be used to prepare food additives; and starch
45 can also be converted into value-added chemical products, such as binders and
46 flocculants (Hanselmann & Burchard, 1996; Wu & Hakkarainen, 2014).

47 However, the modification of starch has been greatly limited by its insolubility in
48 water and most organic solvents. This is especially true if a high degree of substitution
49 (DS) is desired, and in this case modification can only occur in a heterogeneous way.
50 In order to increase the DS, starch esters, for example, were usually prepared in
51 organic solvents, where catalysts may be further added. However, these organic
52 solvents were often volatile, not recyclable, and the catalysts such as pyridine were
53 toxic. These heterogeneous systems have limited the development of starch
54 modification and the application of modified starch (Junistia, Sugih, Manurung,
55 Picchioni, Janssen, & Heeres, 2008).

56 Homogeneous systems have been reported to be beneficial to carbohydrate
57 modification (Heinze & Liebert, 2011; Gao, Luo, & Luo, 2012). As a result, efforts in
58 recent years have been focused on the search for solvents that can completely dissolve
59 starch. Solvents can be classified into derivatizing and non-derivatizing types
60 according to whether or not any chemical reaction is occurring in the dissolution
61 process. Some traditionally- and broadly-used solvents for starch include dimethyl
62 sulfoxide (DMSO) and strong inorganic alkalis (e.g., KOH and NaOH). However,
63 research has shown that alkaline solutions may induce undesirable side reactions such

as structural depolymerization and oxidation (Jackson, Choto-Owen, & Waniska, Rooney, 1988; Jordan, Schmidt, Liebert, & Heinze, 2014). DMSO was found to be able to dissolve starch without structural degradation (Everett & Foster, 1959). However, the dissolution process with DMSO is not straight forward. Starch is heated in 90% aqueous DMSO for about 24 h, followed by precipitation and re-dissolution in water (with further heating); and this procedure requires intensive shear and thermal treatment to starch (Chakraborty, Sahoo, Teraoka, & Gross, 2005; Han & Lim, 2004; Zhong, Yokoyama, Wang, & Shoemaker, 2006).

Ionic liquids (IL's) represent a new class of solvents for starch, which have gained much attention recently. IL's have been claimed as "green solvents" for many substances including natural polymers such as starch and cellulose, because of their chemical and thermal stability, low vapor pressure, and high ionic activity (Gericke, Schlutter, Liebert, Heinze, & Budtova, 2009; Lu, Yan, & Texter, 2009). However, the research on IL's has just been started and the toxicity of IL's remains uncertain. Jastorff et al. and Matsumoto et al. (Jastorff et al., 2003; Matsumoto, Mochiduki, Fukunishi, & Kondo, 2004) have preliminarily studied the toxicity of IL's and proved that the commonly-accepted notion that IL's have low toxicity is incorrect. Most IL's are designed with large organic cations, such as pyridinium, imidazolium and pyrrolidinium while the anions are hexafluorophosphate (PF_6^-), tetrafluoroborate (BF_4^-), dicyanamide ($\text{N}(\text{CN})_2^-$), etc.. Their release to aquatic environments could cause severe water and soil contamination due to the potential toxicity and non-biodegradability (Gathergood & Scammells, 2002). Moreover, the high prices of IL's make them difficult to be applied practically in industry. Therefore, the practical application of IL's remains uncertain.

Another class of solvents for natural polymers is inorganic salt hydrates, which

are liquids that have a designated water-salt mole ratio close to the coordination number of the strongest hydrated ion. Typical examples are liquid phases of solid salt hydrates, such as $\text{Ca}(\text{SCN})_2/\text{H}_2\text{O}$ (Warwicker, Jeffries, Colbran, & Robinson, 1966), $\text{LiSCN}/\text{H}_2\text{O}$ (Lukanoff, Schleicher, & Philipp, 1983) and $\text{ZnCl}_2/\text{H}_2\text{O}$ (Letters, 1932). They are inexpensive, convenient, and environmentally friendly. They are non-derivatizing solvents not only being regarded as very efficient solvents for most natural polymers such as cellulose (Cao, Xu, Chen, Gong, & Chen, 1994), chitin (Wang, Pedersen, Deng, Qiao, & Hou, 2013), but also have been applied as a medium for the chemical functionalization or derivatization, such as carboxymethylation and esterification, of these natural polymers, within a short reaction time (Leipner, Fischer, Brendler, & Voigt, 2000; Sen, Martin, & Argyropoulos, 2013). For example, Heinze et al. used molten $\text{LiClO}_4/3\text{H}_2\text{O}$ as a solvent for homogeneous carboxymethylation of cellulose with a high DS (i.e., <2) in a one-step synthesis after a short reaction time of 4 h (Fischer, Thümmeler, Pfeiffer, Liebert, & Heinze, 2002). Acetylation could also be carried out at a temperature of 130 °C for a short reaction time (0.5-3.0 h) with DS between 1 and 2.5 (Fischer, Leipner, Thümmeler, Brendler, & Peters, 2003). When cellulose is dissolved in a molten salt hydrate, the formation of a homogeneous phase makes cellulose hydroxyl groups completely available for chemical derivatization. This allows the control of both the degree of substitution and the distribution of functional groups, which can undoubtedly broaden the reaction paths and the diversity of products. However, there has been no report on molten salt hydrates applied as solvents or reaction media for starch.

Thus, the aim of the present work was to investigate the ability of ZnCl_2 aqueous solution, a molten salt hydrate, as a solvent for starch, together with the structure and properties of the regenerated starch from this solvent.

2. Materials and Methods

2.1. Materials

Native cassava starch (with moisture content of 14%) was obtained from Dongguan Wenying Starch Trade Co., Ltd., without further treatment. Anhydrous zinc chloride (ZnCl_2) and methanol (analytical grade) were purchased from Guangzhou Chemical Reagent Factory (Guangzhou, China). Other chemicals including hydrazine sulfate and hexamethylenetetramine were of analytical grade and used as received without further purification. All solutions were prepared with distilled water.

2.2. Dissolution and regeneration of starch from zinc chloride solutions

Certain amounts of anhydrous ZnCl_2 were added to a three-neck, round-bottom flask equipped with an electric stirrer to obtain ZnCl_2 aqueous solutions of certain concentrations (65%, 43%, 35% and 29.6%). The ZnCl_2 solution was then mixed with a certain amount of starch (2.0 wt.%, dry weight). Then, the system was stirred at 50 °C for 4 h. In order to prevent degradation by shear force, the revolution speed was set as 60 rpm. Besides, during the dissolution process, the solution was sampled at intervals and observed under a polarized light microscope (Motic BA300POL, Motic Asia, Hongkong) equipped with a digital camera (Moticam Pro 205A, Motic Asia, Hongkong).

Regenerated starch was obtained from the above solutions by adding absolute ethanol with continuous stirring (The absolute ethanol was about 2-5 times the volume of the starch solution). The starch samples were then dried and smashed into powder for further analysis.

ZnCl_2 aqueous solution was acidic (pH = 4.09 for 29.6% and pH = 0.67 for 65%).

For discussing the influence of H^+ on starch, hydrochloric acid (HCl) solution with a pH value of 4.09 was prepared. After that, native cassava starch was treated with the HCl solution under the exactly same processes as with $ZnCl_2$ aqueous solution. And regenerated starch from HCl solution was collected and compared with those from $ZnCl_2$ aqueous solution.

2.3. Characterization of starch and regenerated starch

Fourier-transform infrared (FTIR) spectra for native and regenerated starches were obtained in the range of $400\text{--}4000\text{ cm}^{-1}$ with a TENSOR27 FTIR model manufactured by Burkert, Germany. The samples were mixed with KBr and well ground before being pressed into wafers.

X-ray diffraction (XRD) patterns for native and regenerated starches were obtained on an PW3040/60 X'pert PRO X-ray diffractometer (PANalytical, Netherlands) using the Cu-K α X-ray source operating at 36 kV and 20 mA. The scanning range and rate are $5\text{--}90^\circ$ and $4^\circ/\text{min}$, respectively.

Solid-state ^{13}C CP/MAS spectra were collected on a Bruker Avance 500 nuclear magnetic resonance (NMR) spectrometer using a 4mm MAS BB-1H probe head. The spectral width used was 37.6 KHz. The acquisition time was 27 ms, together with 7.7 μs pre-scan delay and 13.3 μs dwell time. All solid experiments were carried out at 125.8 MHz for ^{13}C with 4096 scans accumulating for each spectrum.

To determine the intrinsic viscosity of native and regenerated starches (Islam, Mohd, & Noor, 2001), the samples were dispersed in 0.5mol/L KOH solutions and stirred in a boiling water bath for 10 min. The solutions were then cooled to room temperature and left overnight. The final concentrations of the solutions were 2.6–6.0 mg/mL. Intrinsic viscosity measurements were obtained using a Ubbelohde

dilution capillary viscometer (size 37, Shanghai Liangjing Glass Instrument Factory, China) immersed in a water bath maintained at 25 ± 0.1 °C. The efflux time of solvent and solutions were measured in triplicate and averaged. Results were expressed as relative viscosity (η_r), $\eta_r = \eta/\eta_0$; specific viscosity (η_{sp}), $\eta_{sp} = \eta_r - 1$; and intrinsic viscosity ($[\eta]$). η is the solution viscosity and η_0 is the solvent viscosity. Intrinsic viscosity, $[\eta]$, was then calculated as the zero concentration-limit of the η_{sp}/c value resulting from the plots of η_{sp}/c versus c straight lines (c represents the concentration of solvent).

The thermal decomposition of native and regenerated starches was determined by a thermogravimetric analysis (TGA) apparatus (TGA4000, PerkinElmer, USA) in an open system (Liu, Yu, Liu, Chen, & Li, 2009). Starch samples were heated to 800 °C at 20 °C/min. The weight loss results were also converted to derivative weight percentages to measure and compare the peak temperatures. The temperature calibration was done by the Curie temperatures of nickel and iron.

3. Results and discussion

3.1. Solubility of starch in $ZnCl_2$ aqueous solutions

Fig. 1 showed the photos of cassava starch (2% concentration) dispersed in $ZnCl_2$ aqueous solution (29.6 wt.%). Starch was seen to be fully dissolved in the solution, as a clear and transparent solution was formed without any sediment (Fig. 1a). In contrast, when starch was dispersed in water, it was only suspended and the solution was turbid (Fig. 1b). In this case, starch granules could only partially swell in water without gelatinization, as the temperature was 50 °C which is lower than the gelatinization temperature of cassava starch.

Since ZnCl_2 aqueous solution was acidic (pH 4.09 at 29.6% concentration), it is necessary to discuss the influence of H^+ on starch. Fig. 1c shows that when starch was dispersed in a HCl solution (pH 4.09), the granules were also suspended in the solvent and no transparent solution could be obtained.

The effect of H^+ on starch granules has been studied since 1885 (Lintner, 1886), and different modified starch products can be produced based on it. Lintnerized starch was the first water soluble starch which was prepared by treatment of starch in 7.5% (w/v) HCl for one week at 22-24 °C, or for three days at 40 °C. Nowadays, commercial soluble starch is still produced by this old procedure (Johnston, Mukerjea, & Robyt, 2011). Besides, microporous starch can also be prepared by treatment of starch in acid solution (Kimura & Robyt, 1995).

While the degradation of starch is inevitable with H^+ in the solution, the process progresses from the amorphous region to the crystalline region (Srichuwong, Isono, Mishima, & Hisamatsu, 2005; Chen, Yu, Simon, Liu, Dean, & Chen, 2011). Specifically, due to the small molecular size, H^+ can penetrate into the pores on the granule surface, which are the radial channels into the granule interior. Since the amorphous region is looser than the crystalline region, H^+ preferably attacks the amorphous region, with the crystalline region being mostly intact. Therefore, the starch granules can only form a classic “Swiss cheese” appearance in acidic solution, and cannot completely be dissolved. In other words, the degradation of starch granules by H^+ can only result in a suspension (as shown in Fig. 1c). Therefore, although ZnCl_2 aqueous solution is acidic, the mechanism for dissolution of starch granules should not be based on the H^+ in the solution.

Fig. 2 showed the birefringence (Maltese crosses) changes of cassava starch granules during dissolution in 29.6% ZnCl_2 aqueous solution. It could be seen that the

starch granules exhibited clear birefringence initially after mixing (Fig. 2a). Then, with the elapse of time, the amount and size of birefringence was observed to decrease (Fig. 2b~2d). After an even longer time (4 h), the visual field gradually turned dark, and all birefringence disappeared totally (Fig. 2d), suggesting that starch granules dissolved completely. Thus, for 2% cassava starch in 29.6% ZnCl_2 aqueous solution at 50 °C, the dissolution time was 4 h.

As is known, the main ions in ZnCl_2 aqueous solution are H^+ , Zn^{2+} and Cl^- . Since starch is slightly acidic in nature and is negatively charged, Cl^- is repelled by starch molecules (Oosten, 1990). H^+ attacked the amorphous region initially, which only resulted in partial degradation of starch granules, but not dissolution. Thus, it was highly possible that the interaction between Zn^{2+} and starch molecules led to the dissolution. Cao et al. (Cao, Xu, Chen, Gong, & Chen, 1994) proved the presence of a zinc-cellulose complex after microcrystalline cellulose (Avicel) was treated with ZnCl_2 aqueous solution. Since starch has the same monosaccharide (glucose) and intermolecular force between molecules (hydrogen bonding) as cellulose, a similar complex may be formed between Zn^{2+} and starch molecules. Furthermore, Islam et al. (Islam, Mohd, & Noor, 2001) found that when starch is added to sodium chloride solution, sodium ions penetrate into starch chains, replacing H^+ which migrates to the water phase; as a result, “sodium salts of starch” are formed. Here in this study, it was possible that a “zinc-starch complex” was formed. Specifically, Zn^{2+} can penetrate into the inner structure of starch, weakening the intra- and inter-molecular hydrogen bonds, disrupting the crystalline region, and forming a “zinc-starch complex” with starch chains. This could be the dominant factor for starch dissolution in ZnCl_2 aqueous solution. Regarding this complexity, further research is required.

Turbidity was also used to determine the dissolution of starch in aqueous ZnCl_2

solution as it represents the cloudiness of a fluid caused by particles in suspension. Turbidity is expected to keep reasonably constant as long as the solute is perfectly soluble, but it will increase sharply with newly added solute remaining solid when the maximum solubility is reached (Mazza, Catana, Garcia, & Cecutti, 2009). As shown in Fig. 3, the starch (2 wt.%) was totally dissolved in 65% ZnCl_2 aqueous solution and formed a clear and transparent solution without any sediment. The turbidity in this case was low (about 21.9 NTU). When the ZnCl_2 concentration was decreased, the turbidity increased very moderately. Nevertheless, when the ZnCl_2 concentration was below 29.6%, precipitation started to appear and the turbidity increased sharply. This suggests that the ZnCl_2 concentration needs to be ≥ 29.6 wt.% for full dissolution of starch (2 wt.%).

3.2. FTIR spectra formative and regenerated starches

The FTIR spectra of native cassava starch and regenerated starch from 29.6% ZnCl_2 aqueous solution were shown in Fig. 4. The spectra of regenerated starch were quite similar to the native counterpart and no new characteristic peaks were found, indicating no chemical changes to the glucosides in the regenerated starch. Thus, the ZnCl_2 aqueous solution can be considered as a non-derivatizing solvent for starch.

Nevertheless, there were also some differences in band wave numbers between native and regenerated starches. As seen from Fig. 4, the O-H bending of adsorbed water at 1647 cm^{-1} was shifted to 1627 cm^{-1} , indicating that the water absorption function of regenerated starch became worse (Vicentini, Dupuy, Leitzelman, Cereda, & Sobral, 2005). Moreover, in the adsorption band between 1100 cm^{-1} and 900 cm^{-1} , which indicates the interactions of hydrogen bonds, the peak at 1016 cm^{-1} was shifted to 1020 cm^{-1} and the one at 929 cm^{-1} was shifted to 935 cm^{-1} . According to Ahmad et

al. and Ma et al. (Ahmad, Anuar, & Yusof, 2011; Ma, Yu, & Zhao, 2006), the peak at the low wave number results from the strong hydrogen bonds that drag the relative groups to a lower vibration frequency, and the peaks at the higher wave number are due to the weak interaction of hydrogen bonding. Thus, the above changes in wave numbers indicated less hydrogen bonding in regenerated starch.

All in all, aqueous ZnCl_2 could be considered as a non-derivatizing solvent for starch without causing obvious change to the starch glucosidic structure but can destroy the inter- and intra-molecular hydrogen bonding in starch.

3.3. Solid state ^{13}C NMR study of native and regenerated starches

For further understanding the effects of ZnCl_2 aqueous solution on starch, ^{13}C NMR spectra on native cassava starch and regenerated starch were obtained (Fig. 5). It can be seen that both native and regenerated starches showed very similar spectra which had four dominant NMR signals. The signals at 90–110 ppm and 81–85 ppm were attributed to C-1 and C-4, respectively, while the signal at 59–65 ppm was assigned to hydroxymethyl C-6. The large signal around 70–79 ppm was associated with C-2, C-3, and C-5 (Morrison, Tester, Gidley, & Karkalas, 1993). Except these, no new peaks appeared in the regenerated starch ^{13}C NMR spectrum.

Compared with the spectra for native and regenerated starch, two distinct differences could be observed: the chemical shifts of C-1 peak (from 103.66 ppm to 105.18 ppm), and the increase in intensity of C-4 peak. The sharp signal intensity of C-4 peak showed an increase in amorphous content (Atichokudomchai, Varavinit, & Chinachoti, 2004; Fan et al., 2013), suggesting that ZnCl_2 aqueous solution destroyed the crystalline region and changed it into amorphous starch. And the shifts of C-1 peak could result from the interaction between Zn^{2+} and hydroxyl groups to form a

complex (Cao, Xu, Chen, Gong, & Chen, 1994). This complex could be the “zinc-starch complex” as mentioned before.

3.4. XRD analysis of native and regenerated starches

Native cassava starch and regenerated starch after treatment with ZnCl_2 aqueous solutions of different concentrations were examined by XRD, and the results were shown in Fig. 6. Native starch exhibited a typical A-type XRD pattern, giving strong reflections at about 15° , 23° and unresolved doublet at nearly 17° and $18^\circ (2\theta)$ (Cheetham & Tao, 1998; Kuo & Lai, 2007). In contrast, the regenerated starch only had a dispersive broad peak and showed no crystalline peaks. This suggested that the crystalline region of native starch was completely damaged during the dissolution. The loss of crystallinity could be attributed the penetration into starch of Zn^{2+} which weakened the inter- and intra- molecular hydrogen bonds of starch (Gao, Luo, & Luo, 2012). These results were in agreement with the shifts in the FTIR and NMR signals.

3.5. Intrinsic viscosity analysis of native starch and regenerated starch

Polymer intrinsic viscosity, $[\eta]$, is an important parameter reflecting the size of a macromolecule in a given solvent at a certain temperature (Liu & Budtova, 2013). This parameter is determined using the classical Huggins equation (see the Materials and Methods section). Fig. 7 shows the intrinsic viscosity of native starch and regenerated starch. Native starch was observed to exhibit the highest intrinsic viscosity (about 192 mL/g). With the increased concentration of ZnCl_2 aqueous solution, the intrinsic viscosity of regenerated starch apparently decreased, from 158 mL/g for 29.6% ZnCl_2 aqueous solution, to 126 mL/g for 65% ZnCl_2 solution.

The intrinsic viscosity of polymers is caused by the friction force between the

solvent and the solute in dilute solution, and can be used to reflect the size of the macromolecular conformation in the certain solution (Wolf, 2007). For starch, no matter amylopectin or amylose, the size of their macromolecular conformation was caused by the degree of polymerization (DP), namely the molecular weight of macromolecules (Higiro, Herald, Alavi, & Bean, 2007). Therefore, from the decreased intrinsic viscosity, it can be deduced that the starch dissolution in ZnCl_2 aqueous solution was accompanied by the degradation of starch chains, which mainly resulted from the attack by H^+ . Since ZnCl_2 aqueous solution of higher concentration had lower pH, the intrinsic viscosity of regenerated starch decreased correspondingly.

Moreover, Fig. 7 also shows the intrinsic viscosity of regenerated starch from a HCl solution (pH 4.09), which was apparently lower than that of regenerated starch from 29.6% ZnCl_2 aqueous solution (pH 4.09). It was possible that the “zinc-starch complex” prevented starch chains from being attacked by H^+ . Nonetheless, this needs further studies. Also, the accurate quantitative relationship between ZnCl_2 solution concentration and starch molecular weight will be discussed in our ongoing research.

3.6. Thermal decomposition of native and regenerated starches

Based on previous research, the thermal decomposition temperature of starch without chemical modification is mainly based on the DP of starch molecular chains, and is not significantly influenced by its crystallinity or granule size (Liu et al., 2013).

The derivative weight percentage of starch samples (DTG) could be seen in Fig. 8. There were two peaks on the DTG curves. The first one represented moisture evaporation, which occurred with the increased temperature until ca. 110 °C. The percentage of weight loss in this step was dependent on the moisture content in starch. The second weight loss peak corresponded to the thermal decomposition of starch

molecular chains. For native starch, this peak located at 337 °C. For regenerated starch from 35%, 43% and 65% ZnCl_2 solutions, this peak located at 271 °C, 253 °C and 250 °C, respectively. The decomposition temperature of native starch was significantly higher than those of regenerated starch, indicating the reduction in DP of starch molecular chains after dissolution in ZnCl_2 solutions. Moreover, the decomposition temperature of regenerated starch was decreased with the increased concentration of ZnCl_2 solution, also showing that a higher concentration of ZnCl_2 solution caused greater degradation of starch macromolecules. These results are in agreement with the intrinsic viscosity results.

4. Conclusions

ZnCl_2 aqueous solution can be used as an effective solvent for starch at 50 °C, which is inexpensive and environmentally friendly. For 2 wt.% starch, the lowest ZnCl_2 concentration to allow full dissolution of starch was 29.6 wt.%, with the dissolving time being 4 h. By characterizing the structure and properties of regenerated starch, ZnCl_2 aqueous solution was found to be a non-derivatizing solvent for starch without causing changes to starch glucosidic structure, but only destroying the inter- and intra- molecular hydrogen bonds of starch and converting the crystalline structure to amorphous state. This is albeit the inevitable degradation of starch chains since ZnCl_2 aqueous solution was acidic. The partial degradation by acid could not be the mechanism of dissolution as it acted mostly to the amorphous region of starch granules. Instead, the dissolution was more likely to rely on the formation of a “zinc-starch complex”.

Further research will be focused on the dissolution mechanism including the verification of the proposed “zinc-starch complex”. Besides, it is worth to study the

interactions between starch and other salt hydrates, the accurate quantitative relationship between ZnCl_2 solution concentration and starch molecular weight, and the chemical modification of starch in a homogeneous phase with ZnCl_2 aqueous solution as a reaction medium.

Acknowledgement

This work is supported by the National Natural Science Foundation of China (NSFC) under the project Nos. 21146011 and 21106023, the Natural Science Foundation of Guangdong Province under the project No. 9151008901000169, the Technology Planning Project of Guangdong Province under the project No. 2013B010403030, and the Science and Technology Projects supported by Guangzhou Education Bureau (Project No. 1201410965).

References

- Ahmad, Z., Anuar, H., & Yusof, Y. (2011). The study of biodegradable thermoplastic sago starch. *Key engineering materials*, 471-472, 397-402.
- Atichokudomchai, N., Varavinit, S., & Chinachoti, P. (2004). A study of ordered structure in acid-modified tapioca starch by ^{13}C CP/MAS solid-state NMR. *Carbohydrate Polymers*, 58, 383-389.
- Cao, N.J., Xu, Q., Chen, C.S., Gong, C.S., & Chen, L.F. (1994). Cellulose hydrolysis using zinc chloride as a solvent and catalyst. *Applied Biochemistry and Biotechnology*, 45, 521-530.
- Chakraborty, S., Sahoo, B., Teraoka, I., & Gross, R.A. (2005). Solution properties of starch nonoparticles in water and DMSO as studied by dynamic light scattering. *Carbohydrate Polymers*, 60, 475-481.

- 388 Cheetham, N.W.H., & Tao, L. (1998). Solid state NMR studies on the structural and
389 conformational properties of natural maize starches. *Carbohydrate*
390 *Polymers*, 36, 285-292.
- 391 Chen, P., Yu, L., Simon, G.P., Liu, X., Dean, K., & Chen, L. (2011). Internal
392 structures and phase-transitions of starch granules during gelatinization.
393 *Carbohydrate Polymers*, 83, 1975-1983.
- 394 Everett, W.W., & Foster, J.F. (1959). The conformation of amylose in solution.
395 *Journal of the American Chemical Society*, 81, 3459-3464.
- 396 Fan, D.M., Ma, W.R., Wang, L.Y., Huang, J., Zhang, F., Zhao, J., Zhang, H., & Chen,
397 W. (2013). Determining the effects of microwave heating on the ordered
398 structures of rice starch by NMR. *Carbohydrate Polymers*, 92, 1395-1401
- 399 Fischer, S., Leipner, H., Thümmeler, K., Brendler, E., & Peters, J. (2003). Inorganic
400 molten salts as solvents for cellulose. *Cellulose*, 10, 227-236.
- 401 Fischer, S., Thümmeler, K., Pfeiffer, K., Liebert, T., & Heinze, T. (2002). Evaluation
402 of molten inorganic salt hydrates as reaction medium for derivatization of
403 cellulose. *Cellulose*, 9, 293-300.
- 404 Gao, J., Luo, Z.G., & Luo, F.X. (2012). Ionic liquids as solvents for dissolution of
405 corn starch and homogeneous synthesis of fatty-acid starch esters without
406 catalysts. *Carbohydrate Polymers*, 89, 1215-1221.
- 407 Gathergood, N., & Scammells, P.J. (2002). Design and preparation of
408 room-temperature ionic liquids containing biodegradable side chains. *Australian*
409 *Journal of Chemistry*, 55, 557-560.
- 410 Gericke, M., Schlüter, K., Liebert, T., Heinze, T., & Budtova, T. (2009). Rheological
411 properties of cellulose/Ionic liquid solutions: From dilute to concentrated states.
412 *Biomacromolecules*, 10, 1188-1194.

- 413 Han, J.A., & Lim, S.T. (2004). Structural changes of corn starches by heating and
414 stirring in DMSO measured by SEC-MALLS-RI system. *Carbohydrate*
415 *Polymers*,55, 265-272.
- 416 Hanselmann, R., & Burchard, W. (1996). Structural properties of fractionated starch
417 polymers and their dependence on the dissolution process. *Macromolecules*, 29,
418 3277-3282.
- 419 Heinze, T., & Liebert, T. (2001). Unconventional methods in cellulose
420 functionalization. *Progress in Polymer Science*, 26, 1689-1762.
- 421 Higiro, J., Herald, T.J., Alavi, S.,& Bean, S. (2007). Rheological study of xanthan and
422 locust bean gum interaction in dilute solution: Effect of salt. *Food Research*
423 *International*,40, 435-447.
- 424 Islam, M.N., Mohd, A.M.D., & Noor, M.A.B.M. (2001). Effect of temperature and
425 starch concentration on theintrinsic viscosity and critical concentration of sago
426 starch (Metroxylon sagu). *Starch/Starke*, 53, 90-94.
- 427 Jackson, D.S., Choto-Owen,C., Waniska, R.D., & Rooney, L.W. (1988).
428 Characterization of starch cooked in alkaliby aqueous high-performance
429 size-exclusion chromatography. *Cereal Chemistry*, 65, 493-496.
- 430 Jastorff, B., Störmann, R.,Ranke, J., Mölter, K., Stock, F., Oberheitmann, B.,
431 Hoffmann, W., Hoffmann, J., Nückter, M., Ondruschka, B., & Filser, J. (2003).
432 How hazardous are ionic liquids? Structure-activity relationships and biological
433 testing as important elements for sustainability evaluation. *Green Chemistry*,5,
434 136-142.
- 435 Johnston, D.A., Mukerjea, R., & Robyt, J.F. (2011). Preparation and characterization
436 of new and improved soluble-starches, -amylose, and -amylopectin by reaction
437 with benzaldehyde/zinc chloride, *Carbohydrate Research*, 346, 2777–2784.

- 438 Jordan, T., Schmidt, S., Liebert, T., & Heinze, T. (2014). Molten imidazole – a starch
439 solvent. *Green Chemistry*, 16, 1967-1973.
- 440 Junistia, L., Sugih, A.K., Manurung, R., Picchioni, F., Janssen, L.P.B.M., & Heeres,
441 H.J. (2008). Synthesis of higher fatty acid starch esters using vinyl laurate and
442 stearate as reactants. *Starch/Starke*, 60, 667-675.
- 443 Kimura, A., & Robyt, J.F. (1995). Reaction of enzymes with starch granules: kinetics
444 and products of the reaction with glucoamylase. *Carbohydrate Research*, 277,
445 87-107.
- 446 Kuo, W.Y., & Lai, H.M. (2007). Changes of property and morphology of cationic
447 corn starches. *Carbohydrate Polymers*, 69, 544-553.
- 448 Leipner, H., Fischer, S., Brendler, E., & Voigt, W. (2000). Structural changes of
449 cellulose dissolved in molten salt hydrates. *Macromolecular Chemistry and*
450 *Physics*, 201, 2041-2049.
- 451 Letters, K. (1932). Viskosimetrische Untersuchungen über die Reaktion von Zellulose
452 mit konzentrierten Chlorzinklösungen. *Kolloidzeitschrift. Collid and Polymer*
453 *Science*, 58, 229-239.
- 454 Lintner, C.J. (1886). Studien über Diastase, *Journal fur praktische chemie-practical*
455 *applications and applied chemistry*. 34, 378–394.
- 456 Liu, W.Q., & Budtova, T. (2013). Dissolution of unmodified waxy starch in ionic
457 liquid and solution rheological properties. *Carbohydrate Polymers*, 93, 199-206.
- 458 Liu, X.X., Wang, Y.F., Yu, L., Tong, Z., Chen, L., Liu, H.S., & Li, X.X. (2013).
459 Thermal degradation and stability of starch under different processing conditions.
460 *Starch/Starke*, 65, 48-60.

- 461 Liu, X.X., Yu, L., Liu, H.S., Chen, L., & Li, L. (2009). Thermal decomposition of
462 corn starch with different amylose/amylopectin ratios in open and sealed systems.
463 *Cereal Chemistry*, 86(4), 383-385.
- 464 Lu, J.M., Yan, F., & Texter, J. (2009). Advanced applications of ionic liquids in
465 polymer science. *Progress in Polymer Science*, 34, 431-448.
- 466 Lukanoff, B., Schleicher, H., & Philipp, B. (1983). Untersuchungen zur Auflösung
467 und Verformung von Cellulose in Schmelzen und konzentrierten Lösungen
468 verschiedener Thiocyanate. *Cellulose Chemistry Technology*, 17, 593-599.
- 469 Ma, X.F., Yu, J.G., & Zhao, A. (2006). Properties of biodegradable poly(propylene
470 carbonate)/starch composites with succinic anhydride. *Composites Science and*
471 *Technology*, 66, 2360-2366.
- 472 Matsumoto, M., Mochiduki, K., Fukunishi, K., & Kondo, K. (2004). Extraction of
473 organic acids using imidazolium-based ionic liquids and their toxicity to
474 *Lactobacillus rhamnosus*. *Separation and Purification Technology*, 40, 97-101.
- 475 Mazza, M., Catana, D.A., Garcia, C.V., & Cecutti, C. (2009). Influence of water on
476 dissolution of selected ionic liquids. *Cellulose*, 16, 207-215.
- 477 Morrison, W.R., Tester, R.F., Gidley, M.J., & Karkalas, J. (1993). Resistance to acid
478 hydrolysis of lipid-complexed amylose and lipid-free amylose in lintnerised
479 waxy and non-waxy barley starches. *Carbohydrate Research*, 245, 289-302.
- 480 Oosten, B.J. (1990). Interactions between starch and electrolytes. *Starch/Stärke*, 42,
481 327-330.
- 482 Sen, S., Martin, J.D., & Argyropoulos, D.S. (2013). Review of cellulose
483 non-derivatizing solvent interactions with emphasis on activity in inorganic
484 molten salt hydrates. *ACS Sustainable Chemistry & Engineering*, 1, 858-870.

- 485 Srichuwong, S., Isono, N., Mishima, T., & Hisamatsu M. (2005). Structure of
486 lintnerized starch is related to X-ray diffraction pattern and susceptibility to acid
487 and enzyme hydrolysis of starch granules. *International Journal of Biological*
488 *Macromolecules*, 37, 115-121.
- 489 Vicentini, N.M., Dupuy, N., Leitzelman, M., Cereda, M.P., & Sobral, P.J.A. (2005).
490 Prediction of cassava starch edible film properties by chemometric analysis of
491 spectra. *Spectroscopy Letters*, 38, 749-767.
- 492 Wang, Y.X., Pedersen, C.M., Deng, T.S., Qiao, Y., & Hou, X.L. (2013). Direct
493 conversion of chitin biomass to 5-hydroxymethylfurfural in concentrated ZnCl_2
494 aqueous solution. *Bioresource Technology*, 143, 384-390.
- 495 Warwicker, J.O., Jeffries, R., Colbran, R.L., & Robinson, R.N. (1966). A Review of
496 the Literature on the Effect of Caustic Soda and other Swelling Agents on the
497 Fine Structure of Cotton. Shirley Institute Pamphlet No. 93, Shirley Institute,
498 Didsbury, Mancheste.
- 499 Wolf, B.A. (2007). Polyelectrolytes revisited: Reliable determination of intrinsic
500 viscosities. *Macromolecular Rapid Communications*, 28(2), 164-170.
- 501 Wu, D., & Hakkarainen, M. (2014). A closed-loop process from microwave-assisted
502 hydrothermal degradation of starch to utilization of the obtained degradation
503 products as starch plasticizers. *ACS Sustainable Chemistry & Engineering*, 2,
504 2172-2181.
- 505 Zhong, F., Yokoyama, W., Wang, Q., & Shoemaker, C.F. (2006). Rice starch,
506 amylopectin, and amylose: molecular weight and solubility in dimethyl
507 sulfoxide-based solvents. *Journal of Agricultural Chemistry*, 54, 2320-2326.
508

1 **Figure Captions**

2 Fig. 1. Pictures of cassava starch (2% concentration) dissolved in 29.6% ZnCl_2
3 aqueous solution (a), water (b) and HCl solution (c) after stirring for 4 h

4 Fig. 2. Birefringence images for 2% cassava starch granules dissolved in 29.6%
5 ZnCl_2 aqueous solution for 0 h (a), 1 h (b), 2 h (c), 3 h (d), and 4 h (e)

6 Fig. 3. Turbidity of starch dissolved in aqueous ZnCl_2 of different concentrations

7 Fig. 4. FTIR spectra for native cassava starch (a) and regenerated starch from
8 29.6% ZnCl_2 aqueous solution (b)

9 Fig. 5. ^{13}C NMR spectra for native starch and regenerated starch from 29.6% ZnCl_2
10 aqueous solution

11 Fig. 6. XRD patterns for regenerated starch samples from 35% (a), 43% (b), 65% (c)
12 ZnCl_2 solutions, and native starch (d)

13 Fig. 7. Intrinsic viscosity of native starch (a), and regenerated starch from 29.6% (b),
14 43% (c), 65% (d) ZnCl_2 solutions, and from HCl solution (pH=4.09) (e)

15 Fig. 8. TGA results for native starch (A), and regenerated starch samples from 35%
16 (B), 43% (C) and 65% (D) ZnCl_2 solution.




Fig. 1. Pictures of cassava starch (2% concentration) dissolved in 29.6% ZnCl_2 aqueous solution (a), water (b) and HCl solution (c) after stirring for 4 h.




Fig. 2. Birefringence images for 2% cassava starch granules dissolved in 29.6% ZnCl_2 aqueous solution for 0 h (a), 1 h(b), 2 h(c), 3 h(d), and 4 h(e).




Fig. 3. Turbidity of starch dissolved in aqueous ZnCl₂ of different concentrations.




Fig. 4. FTIR spectra for native cassava starch (a) and regenerated starch from 29.6% ZnCl₂ aqueous solution (b).




Fig. 5. ^{13}C NMR spectra for native starch and regenerated starch from 29.6% ZnCl_2 aqueous solution.




Fig. 6. XRD patterns for regenerated starch samples from 35% (a), 43%(b), and 65%(c) ZnCl₂ solutions, and native starch (d).




Fig. 7. Intrinsic viscosity of native starch (a), and regenerated starch samples from 29.6% (b), 43% (c), and 65% (d) ZnCl_2 solutions, and HCl solution (pH=4.09) (e).




Fig. 8. TGA results for native starch (A), and regenerated starch samples from 35%(B), 43%(C) and 65% (D) ZnCl_2 solution.